

****FULL TITLE****

*ASP Conference Series, Vol. **VOLUME**, **YEAR OF PUBLICATION***

****NAMES OF EDITORS****

A Photometric Transit Search for Planets around Cool Stars from the Italian Alps: Results from a Feasibility Study

Mario Damasso, Paolo Calcidese, Andrea Bernagozzi and Enzo Bertolini

Astronomical Observatory of the Autonomous Region of the Aosta Valley, Loc. Lignan 39, Nus (Aosta) - Italy

Paolo Giacobbe

Dept. of Physics, University of Turin, Via Giuria 1, I-10125, Turin - Italy

Mario G. Lattanzi, Richard Smart and Alessandro Sozzetti

INAF-Astronomical Observatory of Turin, Strada Osservatorio, 20 I-10025 Pino Torinese (Turin) - Italy

Abstract. A feasibility study was carried out at the Astronomical Observatory of the Autonomous Region of the Aosta Valley demonstrating that it is a well-poised site to conduct an upcoming observing campaign aimed at detecting small-size ($R < R_{\text{Neptune}}$) transiting planets around nearby cool M dwarf stars. Three known transiting planet systems were monitored from May to August 2009 with a 25 cm f/3.8 Maksutov telescope. We reached seeing-independent, best-case photometric RMS less than 0.003 mag for stars with $V \leq 13$, with a median RMS of 0.006 mag for the whole observing period.

1. Introduction

The Astronomical Observatory of the Autonomous Region of the Aosta Valley (OAVdA; 45.7895°N, 7.47833°E) has been identified as a potential site for hosting a photometric transit search for low-mass, small-size planets around nearby cool M dwarf stars. We carried out a study to gauge the near-term and long-term photometric precision achievable, observing known transiting systems in a range of transit depths under variable seeing and sky transparency conditions.

2. Instrumentation and Methodology

We used a 25 cm Maksutov f/3.8 reflector telescope equipped with a CCD Moravian G2-3200ME and an R filter centered at 610 nm (field of view: 52.10 x 35.11 arcmin²; plate scale: 1.43 arcsec/pix; QE $\approx$ 87% at 610 nm). The seeing was monitored each night using a Hartmann mask and the DIMM technique. It varied in the range 1-3 arcsec (median $\approx$ 1.7 arcsec). Data reduction and ensemble differential aperture photometry ($\approx$ 100 reference objects) were carried out with the version 1.0 of an automated, IDL-based pipeline we have developed.




Figure 1. *Left:* WASP-3b transit light curve (28-29 July 2009; expos. 35 s; $T_c=55041.411$) and best-fit curve. *Right:* residuals of the fitted curve.

3. Selected Results

In May-August 2009 we monitored three transiting systems: WASP-3 (Pollacco et al. 2007), HAT-P-7 (Pál et al. 2008) and Gliese 436 (Butler et al. 2004). WASP-3, a $1.24 M_{\text{Sun}}$ star ($V=10.64$) hosting a $1.76 M_{\text{Jupiter}}$ planet (period=1.846834 days), was monitored for 19 nights, often reaching a photometric RMS below 0.003 mag for stars with $V \leq 13$. Fig.1 shows one of the observed WASP-3b transits and the best-fit model is based on the formalism of Mandel & Agol (2002), assuming quadratic limb darkening. We got: $b=0.665 \pm 0.003$ (impact parameter), $i=81.24 \pm 0.06$ degrees (orbital inclination), $r=0.1091 \pm 0.0006$ (planet-to-star radius ratio), $T_c=55041.411 \pm 0.029$ HJD (time of mid transit). The median RMS for the whole observing period is 0.006 mag ($V \leq 13$).

4. Conclusions and Future Work

The feasibility study demonstrated that the OAVdA is a promising site for a long-term photometric survey aimed to detect transiting low-mass, small-size planets. We are planning to set up a system of five 40 cm identical telescopes for an high-precision observing campaign focused on several hundreds nearby M dwarfs, improving the longitudinal coverage of similar programs already ongoing, such as the Arizona-based MEarth project (Nutzman & Charbonneau 2008).

Acknowledgments. M.D., P.C. and A.B. benefit of a grant provided by the European Union, the Autonomous Region of the Aosta Valley and the Italian Department for Work, Health and Pensions.

References

- Butler, P. et al. 2004, ApJ, 617, L580
- Mandel, K. & Agol, E. 2002, ApJ, 580, L171
- Nutzman, P. & Charbonneau, D. 2008, PASP, 120, 317
- Pál, A. et al. 2008 ApJ, 680, 2, 1450
- Pollacco, D. et al. 2007, MNRAS, 385, 1576